

PRIMER NOTE

Characterization of polymorphic microsatellite loci within a young *Boophilus microplus* metapopulation

BROU BASILE KOFFI,* ANGE MARIE RISTERUCCI,† DOMINIQUE JOULIA,* PATRICK DURAND,* NICOLAS BARRÉ,‡ THIERRY DE MEEÛS* and CHRISTINE CHEVILLON*

*UMR CNRS-IRD 2724, Centre IRD, 911 avenue Agropolis, BP 64501, 34 394 Montpellier cedex 5, France, †UMR 1096, CIRAD TA40/03, 34398 Montpellier cedex 5, France, ‡IAC/CIRAD UR 22, BP 73, 98890 Païta, Nouvelle Calédonie, France

Abstract

Nine microsatellite loci from the cattle tick *Boophilus microplus* were isolated and characterized within New Caledonia. This Pacific island was invaded by the cattle tick from a few immigration events dating from mid-20th century. A population survey involving 94 adult ticks indicated monomorphism at one locus, presence of null alleles at another, and high genetic diversity (0.61–0.72) at seven loci apparently suitable for population genetics studies. This opens the opportunity to dissect the populational mechanisms involved in the spectacular capacity of *B. microplus* to cope with local environmental heterogeneity during its recurrent invasions of tropical areas.

Keywords: invasive, pest, ticks

Received 22 November 2005; revision accepted 13 January 2006

Boophilus microplus causes large economic loss in tropical agrosystems because of explosive demography on bovine herds, vector competence for diverse pathogens and recurrent development of acaricide resistances (Frisch 1999; Foil *et al.* 2004; Oliveira *et al.* 2005; and references therein). Another remarkable trait of *B. microplus* has not received attention yet: its ability to quickly adapt to new environmental heterogeneity, as noticed along its short history in New Caledonia. *Boophilus microplus* immigrated there around 1942 (Ragau 1966), before drastic quarantine isolated the island. Since then, the local evolution of this tick has been characterized by invasion of all herds, resistance developments towards all chemicals used in control (Bianchi *et al.* 2003; Ducornez *et al.* 2005) and apparent adaptation to a new invasive host (Barré *et al.* 2001). Neutral markers with rapidly evolving polymorphism are required to analyse the evolutionary determinants of these traits, but published microsatellite markers (Chigagure *et al.* 2000) only lead to null alleles on New Caledonian samples. We thus isolated new microsatellite markers from New Caledonian samples and investigated their potential to characterize polymorphism structure within this isolated and young (*c.* 200 generations old) metapopulation.

DNA was extracted with DNeasy Tissue Kit (QIAGEN) onto a pool (*c.* 1 g) of unfed larvae. Five micrograms of DNA were used to build a microsatellite-enriched library as described in Billotte *et al.* (1999). The enriched microsatellite fragments were amplified by polymerase chain reaction (PCR) into 20 cycles with Rsa21 self-complementary primer (5'-CTCTTGCTTACGCGTGGACTA-3'), cloned into pGEM-T (Promega) and used to transform XL1-Blue MRF supercompetent cells (Stratagene). Randomly picked recombinant colonies lead to 192 inserts that were PCR amplified with Rsa21 primer. PCR products were transferred on Hybond-N+ nylon membranes (Amersham), which were hybridized at 56 °C with [³²P] dATP 5'-end-labelled (GA)₁₅ and (GT)₁₅. Forty-eight positive clones were sent for sequencing (Genome Express). Designing primers with PRIMER 3 (Rozen & Skaletsky 2000; http://www-genome.wi.mit.edu/cgi-bin/primer/primer3_www.cgi), we screened sequences to obtain amplicons sizing between 100 and 220 bp, and retained 25 loci for individual genotyping.

Female blood-meal content was removed to avoid blood PCR-inhibitory effects (Wilson 1997), and individual DNA was extracted with DNeasy Tissue Kit (QIAGEN). Amplifications were performed in 25 µL containing 2.5 µL of 10× buffer (Promega), 10 pmol of each primer, 20 pmol of dNTPs (Invitrogen), 2.5 µL MgCl₂ (25 mM, Promega), 2.5 U

Correspondence: C. Chevillon, Fax: +33 (0)467 41 62 99; E-mail: christine.chevillon@mpl.ird.fr

Table 1 Description of loci (name, GenBank access to reference clone, primers' sequences, fluorescent dye, structure of repeated array), thermocycling conditions (annealing temperature, T_a ; number of cycles, N_c) and genetic variation (alleles number, N_a ; allele size range; observed heterozygosity H_O ; Nei's (1987) average estimate of within-sample gene diversity computed by FSTAT as $H_S = [\bar{n}/(1 - \bar{n})] \cdot [1 - \sum p_{ik}^2 - H_O/2\bar{n}]$ where \bar{n} is the harmonic mean of sample sizes)

Locus name	GenBank Accession no.	Primers (5'-3')	Dye	Repeated array	T_a (°C)	N_c	N_a	Size range (bp)	H_O	H_S
BmA05	DQ001903	F: GCGTGCAGGTTTAGCATA R: ATAGATGAGGGCTTGCCCTTG	PET	(CA) ₅₊₁₀	54	35	3	[110; 114]	0.12	0.27
BmA12	DQ001904	F: CCCCTATAGTGTGCCCTTGA R: CGATGCGTAATACAGGACAA	6-FAM	(CA) ₃₊₇ (CG) ₄	52	35	4	[195; 199]	0.73	0.70
BmA06	DQ001905	F: AATCTGTGCATGGAGGTGTG R: AACCATTAACAAGAAATGCAGTG	NED	(GT) ₈₊₂	54	38	1	101	—	—
BmB12	DQ001906	F: CTGCTCACGGCTGGTTAATTT R: GCACTTTTTTCATCTCCGCTTC	VIC	(TA) ₄ (TG) ₉	51	30	4	[192; 200]	0.69	0.66
BmC03	DQ001907	F: ACGACTTGTGAGGGCGTAACC R: AGATACAGCGCAGCATTTTCG	PET	(CA) ₁₀₊₉	52	35	8	[146; 168]	0.70	0.72
BmC04	DQ001908	F: GAAAAATGTCCGTCGCATT R: CACGAGTCACTCTTCGAGACTTTT	VIC	(TG) ₈	57	30	3	[110; 112]	0.69	0.62
BmC07	DQ001909	F: TTTGACAGGTTTGTGCCATAG R: TCAGCCATATGTTCAACCAGA	NED	(GT) ₁₇	56	35	6	[145; 192]	0.70	0.73
BmD10	DQ001910	F: ACGTGGCGTTTACTGAGTCC R: GGCATATTTTCCACCTCCT	6-FAM	(GT) ₁₃	55	35	5	[151; 155]	0.65	0.65
BmD12	DQ001911	F: CGGCCTCATAATCGTAAAA R: TGGCCACTGGTGACTATGAC	6-FAM	(CA) ₁₀₊₅	50	28	10	[92; 118]	0.65	0.67




Fig. 1 Tick sampling. Cattle and cattle tick are omnipresent on the main island except the south of Noumea. The striped area indicates the highest density of both cattle and cattle tick, and letters indicate the sampling locations. Sample sizes were $N = 24$ in B and C, $N = 22$ in F and $N = 12$ in S and P.

Taq polymerase (Promega) and 2–5 μ L of DNA extracts. Thermocycling consisted in a denaturation step at 94 °C for 3 min followed by 35 cycles of 30 s at 94 °C, 30 s at annealing temperature (T_a), and 30 s at 72 °C. Forward primers were purchased as fluorescently labelled in 5' end (Applied) for use on ABI 310 sequencer. PCR optimization was achievable for nine loci, requiring the thermocycling adjustments detailed in Table 1 and a reduction to 2 μ L $MgCl_2$ (25 mM, Promega) for BmA05 amplification. For a given DNA template, electrophoresis was performed by

pooling 0.5 μ L of internal size standard (GeneScan-500 LIZ, Applied), Hi-Di formamide (20 μ L QSP) and 1 μ L of PCR product of each of three loci differing in fluorescent dye and amplicon size. Allele lengths were measured with GENESCAN (Applied Biosystem).

Consistency in individual genotypic determination among repeated assays was verified, and polymorphism was analysed among five representative samples (Fig. 1). BmA06 was the only monomorphic locus (Table 1). Using MICRO-CHECKER (van Oosterhout *et al.* 2004), we verified the absence of stutter-inducing errors at BmA12, BmC04 and BmD10 loci where single base-pair differences in allele sizes were recorded. MICRO-CHECKER detected heterozygote deficits at BmA05 locus within samples B, C and F, and at BmC04 locus in sample B. These deficits may indicate the presence of null alleles or may result from *B. microplus* biology: development is synchronous within large brotherhoods ($N \geq 100$), eggs are patchily distributed within mother cadavers, neighbour larvae regroup in clusters when seeking an individual host onto develop and mate, and females mate only once. Standard population genetic analyses were performed with FSTAT (Goudet 1995) to settle this point. A low average estimate of gene diversity ($H_S = 0.27$) and a non-null F_{IS} estimate ($F_{IS} = 0.65$, $P < 10^{-4}$) discriminated BmA05 from all other polymorphic loci ($0.62 \leq H_S \leq 0.73$ and acceptance of the hypothesis that ' F_{IS} is not different from 0' with $P > 0.10$). These particularities lead to remove BmA05 polymorphism to investigate further the tick population structure. The

polymorphism recorded at the seven remaining loci involved significant differentiation among samples ($F_{ST} = 0.014$, $P = 0.00005$). Meanwhile, genotypic disequilibrium was detected neither between any two polymorphic loci, nor between sex and either polymorphic locus. Therefore, the seven remaining loci look suitable for population genetics investigation of the local evolutionary determinants of *B. microplus*, at least among New Caledonian and genetically related populations.

Acknowledgements

We thank MM. Abbé, Guillermet, Moglia, Wright and Marcel to have allowed sampling on their cattle, and S. Ducornez for sampling help. This work was funded by the program 'INVABIO2', MEDD, contract n°CV02000234.

References

- Barré N, Bianchi M, Chardonnet L (2001) Role of Rusa deer *Cervus timorensis russa* in the cycle of the cattle tick *Boophilus microplus* in New Caledonia. *Experimental and Applied Acarology*, **25**, 79–96.
- Bianchi MW, Barré N, Messad S (2003) Factors related to cattle infestation level and resistance to acaricides in *Boophilus microplus* tick populations in New Caledonia. *Veterinary Parasitology*, **112**, 75–89.
- Billotte N, Lagoda PJJ, Risterucci AM, Baurens FC (1999) Microsatellite-enriched libraries: applied methodology for the development of SSR markers in tropical crops. *Fruits*, **54**, 277–288.
- Chigagure NN, Baxter GD, Barker SC (2000) Microsatellite loci of the cattle tick *Boophilus microplus* (Acari: Ixodidae). *Experimental and Applied Acarology*, **24**, 951–956.
- Ducornez S, Barré N, Miller RJ, de Garine-Wichatitsky M (2005) Diagnosis of amitraz resistance in *Boophilus microplus* in New Caledonia with the modified Larval Packet Test. *Veterinary Parasitology*, **130**, 285–292.
- Foil LD, Coleman P, Eisler M *et al.* (2004) Factors that influence the prevalence of acaricide resistance and tick-borne diseases. *Veterinary Parasitology*, **125**, 163–181.
- Frisch JE (1999) Towards a permanent solution for controlling cattle tick. *International Journal for Parasitology*, **29**, 57–71.
- Goudet J (1995) FSTAT (version 1.2): a computer program to calculate F-statistics. *Journal of Heredity*, **86**, 485–486.
- Nei M (1987) *Molecular Evolutionary Genetics*, p. 164. Columbia University Press, New York.
- Oliveira MCS, Oliveira-Sequeira TCG, Araujo JP Jr, Amarante AFT, Oliveira HN (2005) *Babesia* spp. infection in *Boophilus microplus* engorged females and eggs in São Paulo State, Brazil. *Veterinary Parasitology*, **130**, 61–67.
- van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004) MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes*, **4**, 535–538.
- Ragau J (1966) Biological observations on ticks (Acari, Agasidae & Ixodidae) in French Pacific Islands. First Symposium for Medical and Veterinary Acarology, Gdansk, 21–26 October.
- Rozen S, Skaletsky HJ (2000) PRIMER 3 on the WWW for general users and for biologist programmers. In: *Bioinformatics Methods and Protocols: Methods in Molecular Biology* (eds Krawetz S, Misener S), pp. 365–386. Humana Press, Totowa, NJ.
- Wilson IG (1997) Inhibition and facilitation of nucleic acid amplification. *Applied and Environmental Microbiology*, **63**, 3741–3751.